

JOB 8: ESTUARINE SEINE SURVEY

JOB 8: ESTUARINE SEINE SURVEY

TABLE OF CONTENTS

	Page
GOAL.....	4
OBJECTIVES	4
METHODS	4
RESULTS	5
Relative Abundance of Juvenile Winter Flounder and Tautog.....	5
Presence of Other Important Recreational Finfish.....	6
Relative Abundance of Forage Species	6
Relative Abundance of Invertebrate Species	7
MODIFICATIONS	7

LIST OF TABLES

Table 8.1: Geometric mean catch of finfish species commonly captured in seine samples, 1988-2019.....	9-10
Table 8.2: Frequency of occurrence of finfish species commonly captured in seine samples, 1988-2019	11-12
Table 8.3: Mean catch of young-of-year winter flounder at eight sites sampled by seine, 1988-2019	13
Table 8.4: Total catch of finfish species commonly captured in seine samples, 1988-2019	14
Table 8.5: Total catch of finfish species infrequently captured in seine samples, 1988-2019	15
Table 8.6: Total catch of invertebrate species captured in seine samples, 2004-2019	16
Table 8.7: Mean catch of forage fish, 1988-2019.....	20

LIST OF FIGURES

Figure 8.1: Sampling locations of the estuarine seine survey along the coast of Connecticut	17
Figure 8.2: Mean catch of all finfish captured in seine samples, 1988-2019	17
Figure 8.3: Mean catch of young-of-year winter flounder, 1988-2019	18
Figure 8.4: Mean catch of young-of-year tautog, 1988-2019	18
Figure 8.5: Mean catch of young-of year black sea bass, 1988-2019.....	19
Figure 8.6: Mean catch of young-of year scup, 1988-2019.....	19
Figure 8.7: Mean catch of forage fish captured in seine samples, 1988-2019.....	20

LIST OF APPENDICES

Appendix 8.1: Finfish species captured in the Estuarine Seine Survey, 1988-2019...	21-22
Appendix 8.2: Invertebrate species captured in the Estuarine Seine Survey, 2019.....	22



Beach seining with 25' bag seine.

JOB 8: ESTUARINE SEINE SURVEY

GOAL

Monitor the abundance and size composition of nearshore young-of-year sportfish and forage fish, along with environmental parameters, to evaluate the effects of fishing and environmental conditions on the distribution and abundance of sportfish in Long Island Sound.

OBJECTIVES

Provide:

1. Indices of abundance for juvenile sportfish (age-0 and 1+).
2. An annual index of forage species abundance.

METHODS

Eight sites (Figure 8.1) are sampled during September using an eight-meter (25 ft.) bag seine with 6.4mm (0.25 in.) bar mesh. Area swept is standardized to 4.6 m (15 ft.), width by means of a taut spreader rope and a 30m (98 ft.), measured distance, parallel to, or at a 45° angle to the shoreline, against the current or tide if present. At each site, six seine hauls are taken within two hours before and after low slack tide during daylight hours. All sites have been sampled since 1988 except Milford which was added in 1990.

Finfish, crabs, and other invertebrates taken in each sample are identified to species or lowest practical taxon (full listing given in Appendix 8.1, 8.2) and counted. One exception is inland silverside, which are not separated from Atlantic silverside because they are rare and difficult to identify. Qualitative counts were used for menhaden when abundant ($n > 1000$) to minimize discard mortality. Winter flounder are measured to total length (mm), and classified as young-of-year (YOY) if less than 12 cm and age 1+ if 12cm or larger. The age of flounder near this size was verified in 1990-1992 by examination of the sagittal otolith. Physical data recorded at each seine location included water temperature and salinity at one-meter depth. The geometric (retransformed natural log) mean catch per standard haul is calculated for total finfish catch and individually for the 22 most abundant species, with separate indices for young-of-year (YOY) and winter flounder age 1 and older. Winter flounder YOY catch is also reported for each site. Confidence intervals (95%) for each geometric mean are retransformations of the corresponding log intervals. Frequency of occurrence is given as a percentage of all samples taken each year.

Diversity in the catch, or species richness, was computed for finfish species captured in the Survey over the time series. Species were divided into three groups based on their temperature preferences and seasonal spawning habits as documented in the literature. Criteria used to assign species into a cold temperate group, warm temperate group, or subtropical group are listed in Job 5.

RESULTS

A total of 48 seine hauls were taken in 2019 at eight sites, yielding a total catch of 13,131 fish of 25 species and 12,655 invertebrates of 14 species. Geometric mean catch of all finfish (fish/haul) in 2019 was 273 fish/haul, above the 30-year time series median of 160 fish/haul (Figure 8.2). Although total catch varies considerably from year to year, the increasing trend is significant ($df=30$, $r^2=0.30$, $p=0.0013$). Dominant species contributing to this increase include young-of-year (YOY) black sea bass, tautog, scup (porgy), northern kingfish, menhaden and a resurgence of five forage species. Other species that contributed to increased total catch in 2019 compared to 2018 were Atlantic Menhaden (85%), Grubby (170%), Inshore Lizardfish (106%), Northern Kingfish (157%), Northern Puffer (98%), Northern Searobin (111%), and Striped Searobin (183%).

Geometric means were calculated for 30 species commonly captured since the survey began in 1988 (Table 8.1). The most frequently caught species was Atlantic silverside, which occurred in all samples, followed by striped killifish, mummichog, tautog, northern pipefish, northern kingfish, northern puffer, scup, black sea bass, and YOY winter flounder (Table 8.2). This rank order has changed from previous years, with a notable increase in forage fish species and winter flounder YOY in 2017 with a decrease in 2018 (Figure 8.3).

Scup, black sea bass and northern kingfish occurrence and abundance increased well above the 30-year time series average in 2016 and 2017 (Tables 8.1 and 8.2). Occurrence of tautog, cunner and sheepshead minnow also ranked high in the time series. Windowpane flounder was again absent in 2019 after re-occurring at low abundance in 2011 and 2014, but absent in 2009-10, 2012-13, and 2015-19 (Table 8.1).

Relative Abundance of Juvenile Winter Flounder and Tautog

The 2019 index of YOY winter flounder (0.688 fish/haul) is 38% greater than the 2018 index. A modest increase from the lowest abundance recorded in 2013 occurred from 2014-2017 (Table 8.3, Figure 8.3). Total catch in 2019 increased 55% from 2018 (Table 8.4). The time series has a significant negative trend ($df=30$, $r^2=0.43$, $p<0.001$), and indicates that a relatively strong year class has not been produced since 1996 (Table 8.1, Figure 8.3). As in previous years, highest abundance was seen at eastern sites (Waterford, Groton,) and Clinton. Bridgeport had no catch (Table 8.3) and the frequency of occurrence of this species has decreased over the time series (Figure 8.3) indicating that juvenile production has contracted in several areas of the Sound. Mean length of YOY winter flounder captured at all sites in 2019 was 69mm and shows no trend over the 30-year time series, ranging from 47.3 to 71.1mm.

The 2019 index of YOY tautog (1.5 fish/haul) was above the series average of 1.0 tautog /haul, increasing from 2017 (1.3 fish/haul) and decreasing from 2015 (4.8 fish/haul) which was the highest abundance in the time series (Table 8.1, Figure 8.4). Overall, the time series has a significant increasing trend ($df=30$, $r^2=0.32$, $p<0.001$). Relatively abundant year classes have been produced in 1998-99, 2002-04, 2007-08, 2012 and 2014-15, 2017-2018.

The frequency of occurrence of this species has also increased over the time series (Figure 8.4) indicating that juvenile production and survival is improving in several areas of the Sound.

Presence of Other Important Recreational Finfish

YOY scup and black sea bass are recent additions to the seine survey (Table 8.1, Figures 8.5 and 8.6). Scup occurred in 1999 but the highest relative abundance has been in the last four years of the time series. Record numbers for YOY scup occurred in 2015 (9.4 fish/haul), with a time series high occurring in 2018 (10.6 fish/haul), which is above the time series mean (1.18 fish/haul).

YOY black sea bass first appeared in Survey catches in 1991 and every year since 1998, reaching their record highest abundance in 2015 (16.3 fish/haul). The 2016 index (5.8 fish/haul) is third highest behind 2014 and 2015. The 2019 index was 2.8 fish/haul, the eighth highest in the times series and slightly lower than 2017 index of abundance. Total catch in 2019 increased 67% from 2018 (Table 8.4).

YOY bluefish show a pattern similar to black seabass, first appearing in the catch in 1991 and almost consistently since 1998. Their abundance increased dramatically in 2014 and 2015, returning to abundance near the time series mean (0.29 fish/haul) in 2016 (Table 8.1). No juvenile bluefish (snappers) were captured in 2017, making it the eighth time in the 30-year time series that no snappers were seen. In 2018, juvenile bluefish reappeared with an abundance of 0.44 fish/haul and increased in 2019 with an abundance of 0.62 fish/haul.

Relative Abundance of Forage Species

Seine survey catches are numerically dominated by forage species, defined here as short-lived, highly fecund species that spend the majority of their life cycle inshore where they are common food items for piscivorous fish. An index of forage fish abundance was generated using the catch of four of the most common forage species caught: Atlantic silverside, striped killifish, mummichog, and sheepshead minnow (Figure 8.7). The 2019 index (161 fish /haul), above the series average of (102 fish/haul) for the time series, increasing from 2017 (107 fish/haul) and decreasing from the 2015 index which was the second highest in the time series.

Although numerically driven by the abundance of silverside, all four forage fish species increased in abundance and occurrence in 2015-2017 and were at or above their time series mean of 102. The time series high occurred in 2003, with a mean catch of 206 forage fish (Table 8.7). Although the forage index varies considerably from year to year, common for short-lived forage species, the increasing trend is significant (r^2 0.16, $p=0.03$, Figure 8.7).

Relative Abundance of Invertebrate Species

A total of 12,655 invertebrates of 14 species or taxon groups were captured in 2019 (Table 8.6, Appendix 8.2), a total similar to 2013 (13,483). Seven crab species were present in the seine hauls, along with two shrimp species. Mud snail, hermit crab, sand shrimp, shore shrimp, green crab, and comb jellyfish were the most abundant and at greater than 50% occurrence (Table 8.6).

Seventy-nine blue crabs were captured in 2019. Blue crabs were captured at Groton, Waterford, Old Lyme, Clinton, New Haven, and Milford sites but continued at relatively low abundances in 2017, 2018 ($n=11$ and 19 crabs) down from a time series high in 2009 ($n=333$ crabs). The 2019 index ($n=79$ blue crabs) is third highest behind 2006 and 2009. The Japanese shore crab re-appeared in 2015 and 2016, with only one captured at the Old Lyme site. No Japanese shore crabs were captured in 2017 or 2018, but one was captured in 2019. The shore shrimp returned to moderate abundance in 2016, 2017, after increasing substantially in 2014-15, then decreasing in 2018 and 2019 (Table 8.6). Sand shrimp decreased significantly from 2015 until 2016, then increased significantly in 2017 until 2019 (Table 8.6). Spider crab abundance has also increased nearly ten-fold from 2011 until 2017 compared to earlier years, with the highest catch in 2017, and decreased significantly in 2018 and 2019 (Table 8.6).

MODIFICATIONS

None.

Beach seining with 25' bag seine at the Old Lyme site.



Table 8.1: Geometric mean catch of finfish species commonly captured in seine samples, 1988-2019.

See Appendix 8.1 for complete taxonomic names.

Species	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
alewife	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
American sand lance	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
American shad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Atlantic menhaden	0.1	0.0	0.0	0.0	0.5	0.0	0.1	0.0	0.0	0.1	0.4	0.4	0.4	0.0	1.0	8.2
Atlantic silverside	68.2	31.6	45.0	88.5	51.2	42.7	37.7	27.0	17.7	23.1	74.3	102.5	99.7	36.1	80.1	113.6
Atlantic tomcod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
black sea bass	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.1	0.0	0.0	0.1	0.1	0.0	1.0	0.4	0.2
blueback herring	0.0	0.1	0.0	0.5	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.1
bluefish	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.9	0.0	0.1	0.0	0.2
cunner	0.2	0.3	0.0	0.1	0.2	0.0	0.3	0.2	0.3	0.0	0.3	0.5	0.3	0.2	0.3	0.2
fourspine stickleback	0.3	0.4	0.0	0.7	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.1	0.0	0.0
grubby	0.8	0.1	0.0	0.1	0.5	0.1	0.4	0.3	0.2	0.3	0.2	0.5	0.1	0.2	0.3	0.5
inshore lizardfish	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.4	0.1	0.2	0.2	1.2	0.0	0.0
mummichog	2.8	1.6	1.1	1.9	1.6	3.7	3.3	0.7	1.2	0.5	2.0	0.8	3.2	1.4	3.4	2.9
naked goby	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
northern kingfish	0.0	0.0	0.0	0.0	0.1	0.2	0.0	0.1	0.0	0.1	0.1	0.1	0.0	0.2	0.1	0.2
northern pipefish	0.7	0.3	0.4	1.0	0.9	0.9	1.1	0.5	1.0	0.4	2.1	1.0	1.0	1.4	0.5	0.3
northern puffer	0.1	0.3	0.1	0.4	0.1	0.4	0.2	0.5	0.2	0.1	0.1	0.2	0.6	0.2	0.7	0.7
rainbow smelt	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
scup	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.0	0.6
sheepshead minnow	0.8	1.0	0.1	0.6	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.4	0.2	0.6	0.7
smallmouth flounder	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.3	0.0	0.1	0.0	0.0
striped bass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
striped killifish	11.9	7.9	5.9	4.2	3.1	4.9	5.1	3.9	2.0	1.5	7.2	4.5	8.6	7.5	14.5	14.9
striped searobin	0.2	0.2	0.1	0.2	0.1	0.9	0.1	0.0	0.1	0.4	1.9	0.6	0.1	0.4	0.3	0.7
summer flounder	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
tautog	0.3	0.1	0.3	0.7	0.4	0.2	0.8	0.7	0.3	0.2	0.9	1.3	0.5	0.6	1.5	1.1
weakfish	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
windowpane flounder	0.6	0.1	0.2	0.2	0.3	0.3	0.1	0.2	0.7	0.4	0.1	0.1	0.1	0.0	0.0	0.1
winter flounder-age 1+	0.2	0.1	0.0	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.0	0.0	0.0
winter flounder YOY	15.4	1.7	2.9	5.2	11.9	5.7	14.2	10.1	19.2	7.5	9.2	8.7	4.3	1.3	3.1	8.1

Table 8.1 continued: Geometric mean catch of finfish species commonly captured in seine samples, 1988-2019.

See Appendix 8.1 for complete taxonomic names.

Species	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
alewife	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
American sand lance	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
American shad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.3	0.0
Atlantic menhaden	0.4	0.2	0.4	0.6	0.1	0.3	0.0	0.1	0.0	0.1	1.2	9.9	0.4	0.5	1.6	1.4
Atlantic silverside	85.1	81.3	37.7	74.9	57.5	66.8	96.9	66.5	45.0	35.0	64.8	114.5	73.0	75.9	117.4	111.3
Atlantic tomcod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
black sea bass	0.4	0.1	0.5	0.6	0.3	1.1	0.4	3.2	5.2	3.7	10.8	16.3	5.8	3.2	4.6	2.8
blueback herring	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.8	0.0
bluefish	0.2	0.1	0.2	0.0	0.0	0.3	0.0	0.2	0.4	0.1	0.8	3.4	0.3	0.0	0.4	0.6
cunner	0.5	0.3	0.1	0.5	0.1	0.2	0.1	0.0	0.4	0.0	0.5	0.1	0.0	0.2	0.2	0.2
fourspine stickleback	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0
grubby	1.3	0.8	0.3	0.3	0.2	0.5	0.3	0.7	0.2	0.2	0.2	0.2	0.0	0.3	0.1	0.6
inshore lizardfish	0.0	0.0	1.9	0.2	0.3	0.2	0.1	0.2	0.2	0.1	1.6	0.4	0.0	0.4	0.3	0.6
mumichog	2.3	1.5	2.5	7.3	2.9	3.8	1.7	3.1	1.6	0.9	5.0	5.3	2.2	3.3	4.5	5.6
naked goby	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0
northern kingfish	0.3	0.1	0.0	0.0	0.2	0.3	0.5	0.2	0.5	0.7	1.1	1.0	0.1	0.8	0.3	2.0
northern pipefish	0.7	0.5	0.6	0.8	0.7	1.9	0.6	1.1	1.4	1.7	2.6	2.0	0.5	0.8	2.2	1.4
northern puffer	0.7	0.5	0.4	1.2	0.2	0.3	0.4	0.4	0.9	1.1	1.1	1.4	0.2	0.2	0.6	1.5
rainbow smelt	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
scup	0.2	0.9	0.1	1.0	0.1	1.9	0.1	0.2	2.1	0.1	2.6	9.4	1.3	2.5	10.6	2.7
sheepshead minnow	0.5	0.2	0.2	3.3	1.2	0.5	0.3	0.5	0.8	0.2	0.6	0.3	0.5	0.5	1.8	1.0
smallmouth flounder	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.9	0.4	0.5	0.1	0.2	0.1	0.0	0.1	0.0
striped bass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
striped killifish	12.9	19.4	7.1	21.2	21.7	12.3	15.9	28.7	5.3	3.8	14.5	17.1	10.2	10.4	23.7	20.2
striped searobin	0.5	0.2	0.1	0.3	0.3	0.8	0.2	0.1	0.1	0.2	1.1	0.7	0.0	0.0	0.0	0.4
summer flounder	0.0	0.0	0.2	0.1	0.1	0.0	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0
tautog	1.4	0.7	0.4	2.4	1.0	0.4	0.4	0.3	1.3	0.6	3.5	4.8	1.1	1.3	3.7	1.5
weakfish	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
windowpane flounder	0.2	0.2	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
winter flounder	0.1	0.2	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
winter flounder YOY	11.0	5.6	0.9	4.7	2.0	0.8	1.0	1.1	0.3	0.3	0.5	0.6	0.6	1.0	0.4	0.7

Table 8.2: Frequency of occurrence of finfish species commonly captured in seine samples, 1988-2019.

See Appendix 8.1 for complete taxonomic names.

Species	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
alewife	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.02
American sand lance	0.00	0.00	0.00	0.00	0.02	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
American shad	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Atlantic menhaden	0.06	0.05	0.04	0.04	0.19	0.06	0.10	0.04	0.00	0.06	0.06	0.15	0.10	0.02	0.27	0.58
Atlantic silverside	0.97	0.93	0.96	1.00	1.00	0.96	1.00	0.96	0.94	0.92	0.98	0.94	1.00	0.92	1.00	0.96
Atlantic tomcod	0.00	0.00	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00
black sea bass	0.00	0.00	0.00	0.04	0.00	0.00	0.15	0.04	0.00	0.00	0.06	0.08	0.02	0.25	0.17	0.13
blueback herring	0.00	0.05	0.04	0.13	0.04	0.00	0.06	0.02	0.00	0.00	0.02	0.08	0.02	0.00	0.04	0.06
bluefish	0.00	0.00	0.00	0.10	0.02	0.00	0.02	0.00	0.00	0.02	0.13	0.46	0.04	0.13	0.02	0.10
cunner	0.17	0.19	0.04	0.10	0.15	0.00	0.23	0.15	0.13	0.02	0.21	0.23	0.19	0.15	0.13	0.17
fourspine stickleback	0.17	0.19	0.00	0.23	0.15	0.04	0.02	0.00	0.04	0.00	0.13	0.04	0.02	0.06	0.00	0.00
grubby	0.33	0.07	0.04	0.10	0.31	0.06	0.33	0.25	0.19	0.29	0.17	0.27	0.10	0.17	0.21	0.29
inshore lizardfish	0.06	0.00	0.04	0.00	0.00	0.06	0.10	0.00	0.00	0.29	0.06	0.17	0.19	0.56	0.04	0.00
mummichog	0.47	0.48	0.35	0.40	0.38	0.50	0.42	0.35	0.42	0.15	0.42	0.29	0.44	0.42	0.54	0.44
naked goby	0.00	0.00	0.02	0.06	0.00	0.00	0.00	0.02	0.00	0.00	0.02	0.02	0.00	0.08	0.02	0.02
northern kingfish	0.00	0.00	0.00	0.06	0.08	0.10	0.04	0.15	0.04	0.13	0.10	0.08	0.04	0.13	0.04	0.15
northern pipefish	0.42	0.31	0.37	0.63	0.35	0.50	0.58	0.33	0.44	0.33	0.73	0.48	0.54	0.48	0.19	0.25
northern puffer	0.08	0.24	0.09	0.27	0.08	0.31	0.17	0.40	0.15	0.06	0.10	0.19	0.35	0.17	0.35	0.31
rainbow smelt	0.00	0.00	0.00	0.00	0.00	0.02	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
scup	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.23	0.35	0.25
sheepshead minnow	0.31	0.31	0.09	0.21	0.04	0.02	0.02	0.04	0.00	0.04	0.04	0.06	0.17	0.10	0.15	0.19
smallmouth flounder	0.03	0.00	0.00	0.02	0.00	0.13	0.10	0.06	0.04	0.04	0.00	0.21	0.06	0.13	0.00	0.00
striped bass	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.06
striped killifish	0.78	0.67	0.65	0.73	0.58	0.65	0.58	0.69	0.54	0.40	0.75	0.67	0.63	0.71	0.85	0.81
striped searobin	0.11	0.12	0.11	0.10	0.08	0.48	0.10	0.02	0.10	0.35	0.60	0.38	0.10	0.29	0.25	0.40
summer flounder	0.00	0.00	0.00	0.00	0.00	0.04	0.10	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00
tautog	0.22	0.05	0.22	0.42	0.31	0.19	0.33	0.33	0.13	0.17	0.38	0.46	0.23	0.40	0.54	0.50
weakfish	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.13
windowpane flounder	0.31	0.10	0.13	0.23	0.23	0.19	0.17	0.19	0.35	0.23	0.13	0.13	0.06	0.00	0.02	0.10
winter flounder –age 1+	0.25	0.12	0.00	0.15	0.08	0.23	0.17	0.19	0.10	0.15	0.10	0.06	0.15	0.04	0.02	0.00
winter flounder YOY	0.97	0.71	0.74	0.92	0.98	0.88	0.98	0.94	1.00	0.94	0.92	0.88	0.77	0.58	0.79	0.85

Table 8.2 cont.: Frequency of occurrence of finfish species commonly captured in seine samples, 1988-2019. See Appendix 8.1 for complete taxonomic names.

Species	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
alewife	0.00	0.00	0.00	0.00	0.00	0.00	0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02
American sand lance	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
American shad	0.00	0.00	0.00	0.00	0.00	0.00	0	0.00	0.00	0.00	0.00	0.04	0.00	0.13	0.1	0.00
Atlantic menhaden	0.08	0.06	0.13	0.17	0.02	0.15	0.02	0.02	0.04	0.04	0.23	0.54	0.1	100	0.29	0.23
Atlantic silverside	1	1	1	1	1	1	1	1	0.98	1	4	1	1	0.00	1	1
Atlantic tomcod	0.02	0.02	0.00	0.00	0.02	0.00	0	0.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02
black sea bass	0.25	0.08	0.23	0.23	0.15	0.27	0.13	0.58	0.75	0.58	0.77	0.9	0.88	0.77	0.94	0.50
blueback herring	0.00	0.00	0.00	0.02	0.00	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.23
bluefish	0.15	0.04	0.08	0.00	0.02	0.15	0.02	0.1	0.21	0.08	0.23	0.77	0.21	0.00	0.27	0.35
cunner	0.29	0.21	0.13	0.25	0.1	0.17	0.08	0.04	0.23	0.02	0.31	0.1	0.00	0.13	0.13	0.15
fourspine stickleback	0.02	0.00	0.02	0.00	0.00	0.02	0	0.04	0.00	0.00	0.15	0.00	0.04	0.06	0.08	0.00
grubby	0.5	0.46	0.27	0.15	0.19	0.27	0.21	0.42	0.23	0.2	0.19	0.15	0.02	0.19	0.08	0.40
inshore lizardfish	0.06	0.00	0.6	0.13	0.19	0.15	0.13	0.1	0.15	0.13	0.6	0.25	0.00	0.29	0.29	0.31
mummichog	0.35	0.27	0.48	0.65	0.48	0.5	0.4	0.42	0.35	0.27	0.54	0.65	0.4	0.5	0.54	0.65
naked goby	0.04	0.00	0.08	0.00	0.02	0.00	0	0.02	0.08	0.06	0.08	0.02	0.02	0.04	0.04	0.00
northern kingfish	0.17	0.1	0.02	0.02	0.19	0.17	0.23	0.13	0.29	0.35	0.4	0.38	0.1	0.4	0.21	0.56
northern pipefish	0.48	0.25	0.29	0.42	0.23	0.52	0.4	0.44	0.6	0.6	0.69	0.75	0.31	0.5	0.83	0.58
northern puffer	0.4	0.31	0.29	0.44	0.23	0.23	0.21	0.31	0.42	0.38	0.48	0.31	0.21	0.23	0.38	0.54
rainbow smelt	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
scup	0.13	0.29	0.04	0.29	0.02	0.38	0.04	0.06	0.42	0.08	0.48	0.71	0.38	0.71	0.96	0.54
sheepshead minnow	0.15	0.15	0.06	0.4	0.27	0.13	0.1	0.13	0.25	0.07	0.17	0.13	0.13	0.15	0.4	0.23
smallmouth flounder	0.00	0.00	0.02	0.00	0.13	0.15	0.06	0.4	0.17	0.29	0.06	0.15	0.13	0.02	0.1	0.02
striped bass	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02
striped killifish	0.73	0.96	0.65	0.88	0.94	0.75	0.9	0.98	0.65	0.58	0.88	0.88	0.79	0.85	1	0.94
striped searobin	0.38	0.13	0.13	0.27	0.19	0.4	0.17	0.06	0.08	0.15	0.49	0.29	0.02	0.04	0.04	0.21
summer flounder	0.00	0.00	0.19	0.06	0.15	0.02	0.04	0.00	0.08	0.12	0.06	0.13	0.02	0.02	0.04	0.04
tautog	0.54	0.42	0.17	0.54	0.42	0.35	0.31	0.23	0.6	0.33	0.63	0.83	0.67	0.5	0.85	0.65
weakfish	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.08	0.00
windowpane flounder	0.21	0.15	0.06	0.04	0.1	0.00	0.04	0.02	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00
winter flounder –age 1+	0.17	0.21	0.15	0.08	0.15	0.04	0.04	0.04	0.05	0.00	0.06	0.04	0.02	0.06	0.02	0.04
winter flounder YOY	0.98	0.94	0.46	0.92	0.71	0.52	0.6	0.63	0.27	0.23	0.33	0.46	0.35	0.48	0.4	0.46

Table 8.3: Mean catch of young-of-year winter flounder at eight sites sampled by seine, 1988-2019.

BPT=Bridgeport, CLT=Clinton, GRT=Groton, GRW=Greenwich, MIL=Milford, OLM=Old Lyme, WTF=Waterford

Year	BPT	CLT	GRT	GRW	MIL	NHH	OLM	WTF	All Sites
1988	*18.72	2.73	11.39	9.63	-	38.66	58.19	29.57	15.4
1989	1.70	1.14	1.53	0.70	-	2.14	2.04	2.99	1.7
1990	3.97	0.19	2.21	0.51	1.62	5.69	16.83	2.64	2.9
1991	1.77	4.10	5.62	1.99	2.46	6.45	15.32	18.25	5.2
1992	3.34	5.53	6.25	9.42	4.29	40.15	47.99	32.52	11.9
1993	1.22	1.40	8.59	4.33	3.62	11.47	13.34	16.66	5.7
1994	4.46	8.11	38.36	4.26	4.62	35.34	61.65	21.03	14.2
1995	1.94	3.19	30.28	7.22	1.77	18.93	34.23	36.58	10.1
1996	7.67	11.81	15.67	*12.61	*6.58	*49.29	91.34	30.53	*19.2
1997	2.87	6.61	23.69	3.43	1.64	3.79	52.01	11.25	7.5
1998	1.24	4.03	17.63	8.12	0.91	22.37	57.19	21.89	9.2
1999	1.04	2.60	25.7	7.95	3.49	0.94	*137.07	36.12	8.7
2000	2.14	0.51	0.76	6.65	0.78	1.74	48.34	*41.56	4.3
2001	0.20	1.12	4.12	1.24	0.59	0	0.91	9.10	1.3
2002	0.91	2.66	3.06	5.08	0.26	1.08	15.55	8.98	3.1
2003	1.88	4.61	*45.78	5.88	0.89	1.70	51.13	32.30	8.1
2004	1.00	*18.36	33.84	11.27	3.36	33.06	11.13	13.04	11.0
2005	1.94	11.14	16.7	7.71	5.14	1.64	4.06	7.30	5.6
2006	0.12	1.38	5.53	0.12	0	0	3.30	1.29	0.9
2007	0.78	5.65	17.90	4.44	0.78	6.42	7.89	7.11	4.7
2008	0.51	2.45	10.84	0.51	0	1.57	2.62	5.94	2.0
2009	0.91	1.62	2.29	0.12	0.51	0.12	0.12	1.75	0.8
2010	0.41	1.11	1.71	1.33	0.12	0.41	1.88	1.57	1.0
2011	0.12	0.98	1.18	2.26	0.78	0.12	4.27	1.45	1.1
2012	0	0.26	0.70	0.76	0	0.12	0.26	0.44	0.3
2013	0	0	1.14	0.26	0	0	0.65	0.57	**0.28
2014	0.12	0.12	1.82	0.26	0.12	0.12	1.35	0.65	0.47
2015	0	0.59	1.96	0.70	0.12	0.12	0.51	2.40	0.64
2016	0.12	0	1.49	0.20	0	0	1.14	6.03	0.63
2017	0	0.12	13.53	0.91	0.26	0.26	0.12	4.19	1.03
2018	0	0.31	1.29	0.41	0	0	0.41	1.80	0.42
2019	0	0.65	2.56	0.44	0.26	0.12	0.41	2.91	0.68

*record high for a site/year. **record low for time-series

Table 8.4: Total catch of finfish species commonly captured in seine samples, 1988-2019. See Appendix 8.1 for complete taxonomic names.

Species	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	Total
Atlantic Menhaden	3	2	2	4	1,074	3	9	2		11	2,003	377	1,236	1	1,284	5,098	1,117	75	117	144	21	54	3	43	2	14	3,404	3,948	150	703	614	1514	23,032
Atlantic Silverside	4,750	3,316	5,356	6,383	5,468	5,263	6,311	2,352	1,942	3,249	6,345	10,120	8,738	4,417	5,730	13,278	5,122	5,089	3,267	5,087	3,245	4,156	7,063	4,657	4,142	3,958	3,832	7,549	6,459	4,869	6,963	6,999	175,475
Black Sea Bass				10			41	43		27	14	2	687	63	27	110	15	82	109	33	304	86	489	783	1,197	1,950	1,794	500	338	267	537	9,508	
Bluefish			3	194	10		5	2		3	24	1	13	5	23	8	30		7	53	1	26	54	17	194	289	45	44	55	1106			
Cunner	15	27	2	5	19		42	24	63	1	23	142	26	15	110	15	54	35	18	58	8	28	15	2	42	1	73	7		28	23	15	936
Fourspine Stickleback	33	76		183	11	21	1		3	24	3	1	7		9				2		8		2			13		2	3	11	413		
Grubby	111	3	2	7	61	6	38	19	21	28	17	55	15	73	33	95	143	76	31	32	16	51	25	55	18	19	18	16	1	35	4	50	1,174
Inshore Lizardfish	5		2			4	6			46	6	16	15	103	2	3		169	18	26	22	10	16	23	11	135	38		30	19	62	787	
Mummichog	1,031	197	171	765	573	1,256	1,943	78	149	190	396	115	1,008	246	811	702	637	543	398	1,203	498	857	299	775	329	199	1,098	999	519	676	748	791	20,200
Naked Goby				1	4				1		1	1		4	2	2	2		13		2		2	4	4	6	5	1	7	2	64		
Northern Kingfish				3	4	23	2	9	3	10	7	6	5	17	5	21	38	11	1	1	23	42	76	30	54	81	149	113	10	142	24	199	1,109
Northern Pipefish	65	23	33	106	120	82	117	52	241	38	295	141	96	189	87	25	72	92	82	75	156	307	49	248	152	204	413	142	48	63	147	126	4,086
Northern Puffer	4	22	13	34	4	37	15	40	25	5	5	13	63	14	79	101	75	93	34	241	19	41	51	28	98	202	97	448	18	17	56	163	2,155
Northern Searobin	2	1				1	1				3	40	24	5	4	13	2	10			1	9		6	35	105		45	41	143	491		
Rainwater Killifish									3	4			2		6	35	53	19	3								4	55	6	190			
Scup											1		58	172	131	50	154	6	170	14	413	21	30	375	18	485	1,573	198	212	646	532	5,259	
Sheepshead Minnow	174	815	5	345	4	1	2	30		14	19	12	267	59	402	276	205	28	104	1,439	304	203	82	219	238	59	154	60	742	352	296	340	7,250
Smallmouth Flounder	1			1		8	14	7	2	5		40	3	12					1		14	21	5	114	63	49	15	13	7	1	10	1	407
Striped Killifish	1,511	1,383	748	659	465	773	1,923	520	269	289	1,066	539	1,797	1,494	1,698	3,410	1,548	1,470	1,063	1,994	1,874	1,508	1,300	1,964	720	493	1,158	1,531	1,482	961	1,412	1,358	40,380
Striped Searobin	22	12	5	94	5	71	5	1	9	40	178	51	7	33	33	62	38	19	6	32	36	82	14	4	7	14	121	84	1	2	2	45	1,135
Summer Flounder						2	6		1		1								16	8	8	1	6		6	7	3	11	1	2	2	2	83
Tautog	23	5	23	72	32	16	104	88	42	19	135	174	67	59	153	140	145	64	93	321	131	25	33	27	123	73	467	446	75	140	255	141	3,711
Windowpane Flounder	49	4	22	19	35	30	9	13	71	50	12	10	4		1	5	15	15	3	2	17		2	4								394	
Winter Flounder 1+	12	6		7	6	14	13	12	21	282	9	4	7	2	3		9	11	7	6	13	2	2	2	2		3	2	1	3	1	2	464
Winter Flounder YOY	900	117	276	410	1,055	483	1,401	916	1,486	874	999	1,497	708	138	302	1,310	914	470	110	365	190	72	71	86	22	24	48	48	74	140	29	51	15,586

Table 8.5: Total catch of finfish species infrequently captured in seine samples, 1988-2019. See Appendix 8.1 for complete taxonomic names

Species	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	Total
Alewife								1							28	1															1	31	
American Eel	1	3						1				5																		2	1	31	
American Sand Lance				1			10																									24	
American Shad	18	1									151																			42	50	262	
Anchovy, spp (YOY)																					15										3,051	3,066	
Atlantic Needlefish																	2													1		3	
Atlantic Tomcod						3					1					1	3				1									2	19		
Banded Butterflyfish											2	3				4	2	3	1	3										3	3		
Banded Gunnel																																1	19
Banded Rudderfish																					1											1	
Bay Anchovy								4	69		27		1	11		1	12					1					520	24			670		
Black Spotted Stickleback																															2	2	
Blue Spotted Coronet Fish											1																				2	3	
Blueback Herring	26																			9		3	1	1						11	165	216	
Burrfish, Striped		15	2		1			1	9	142	3	8	2	17								10		4									214
Butterfish											1																					22	
Crevalle Jack	6						1															1										8	
Feather Blenny																						36									7	43	
Flying Gurnard																	1															1	
Gizzard Shad																				4											4		
Grey Snapper		1																														1	
Hogchoker				2																1											3		
Lined Seahorse			4					1			2									2	7	2	1	2					1	4	7	33	
Little Skate						1					1			1																	2		
Moonfish																				1											16	16	
Northern Sennet																					1											1	
Northern Star Gazer		5																														5	
Oyster Toadfish	5			1							1	1			1	1	2	1	1	1	2	1					6	2	4	2	3	39	
Pumpkin Seed			2														3															5	
Rainbow Smelt					5	2											34															41	
Rock Gunnel		1		1	1	1					3						1					1										9	
Smooth Dogfish		1																															1
Spot																																6	
Striped Anchovy																				3												6	
Striped Bass																																1	
Threespine Stickleback															11																	11	
Weakfish																15															4		
Web Burrfish																	1														10		
White Mullet	1	1	8		3										1				7	7	11		75	68	22				15	31	16	266	
White Perch																	3			11				6									20
Yellow Jack																				1													1

Table 8.6: Total catch of invertebrate species captured in seine samples, 2004-2019.

See Appendix 8.2 for complete taxonomic names.

Species	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
Blue crab	1	2	84	31	4	333	35	23	27	18	17	18	6	11	19	79	
Boreal squid				1													
Brown shrimp			11										3				
Channeled whelk							1				3						
Common slipper shell			13														
Flat-claw hermit crab	761	532	703	153	244	539	558	441	283	367	562	308	2,878	829	658	1,157	
Green crab	234	266	341	147	644	176	308	228	175	253	273	213	256	460	432	345	
Horseshoe crab													1				
Japanese shore crab	1		1	1				6	1			1	1			1	
Jonah crab						2											
Knobbed whelk																1	
Lady crab	298	119	66	195	92	42	19	24	18	13	41	102	12	20	2	52	
Mantis shrimp										1							
Mole crab	1	5															
Moon jellyfish						319							461	16	200	200	
Mud crab	60	55	74	30	85	67	308	80	80	1,100	43	142	9	1	20	31	
Mud snail	948	2,071	4,478	3,569	3,810	3,128	2,699	2,683	3,072	5,787	6,938	11,132	11,687	6,061	6,927	9,422	
Northern comb jelly						346	36			3,620	1,200			185	1,648	111	300
Northern moon snail														6		2	
Oyster drill			38													69	
Rock crab	2						1										
Sand shrimp	278	373	1,027	525	2,625	762	902	1,507	246	1,794	662	207	33	518	650	531	
Scallop (bay)												3	3	1	2		
Shore shrimp	990	404	1,149	707	1,390	535	619	762	402	511	1,011	4,795	478	1,517	921	526	
Spider crab	4	5	6	1	3	1	7	33	13	20	14	45	53	161	21	8	
Squid (longfin)												6					
Starfish spp.							1										




Figure 8.1: Sampling locations of the Estuarine Seine Survey.




Figure 8.2: Mean catch of all finfish captured in seine samples, 1988-2019. Geometric mean catch (numbers) per haul includes samples at all sites. Note that sampling at the Milford site began in 1990.

Winter Flounder YOY Geometric Mean Catch, 1988-2019




Figure 8.3: Mean catch and occurrence of young-of-year winter flounder, 1988-2019. The negative trend (dashed line) is significant ($r^2 = 0.43$, $p < 0.001$, $df = 30$).

Tautog Geometric Mean Catch, 1988-2019




Figure 8.4: Mean catch and occurrence of young-of-year tautog, 1988-2019. The positive trend (dashed line) is significant ($r^2 = 0.32$, $p < 0.001$, $df = 30$).

Black Sea Bass Geometric Mean Catch, 1988-2019




Figure 8.5: Mean catch of black seabass young-of-year, 1988-2019. The positive trend (dashed line) is significant ($r^2 = 0.38$, $p < 0.001$, $df = 30$).

Scup Geometric Mean Catch, 1988-2019




Figure 8.6: Mean catch of scup young-of-year, 1988-2019. Annual percent of hauls catching one or more scup are also shown. The positive trend (dashed line) is significant ($r^2 = 0.33$, $p < 0.0005$, $df = 30$).




Figure 8.7: Mean catch of forage fish, 1988-2019. Forage species included in the index are Atlantic silversides, striped killifish, mummichog, and sheepshead minnow. See Appendix 8.1 for complete taxonomic names.

Table 8.7: Mean catch of forage fish, 1988-2019.

1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	1999
139	62	64	110	71	65	57	42	26	32	100	127	
2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010		
146	52	125	206	130	122	59	149	100	106	137		
2011	2012	2013	2014	2015	2016	2017	2018	2019				
127	60	46	104	171	99	107	167	161				

Appendix 8.1: Finfish species captured in the Estuarine Seine Survey, 1988-2019.

<u>COMMON NAME</u>	<u>SPECIES CODE</u>	<u>SCIENTIFIC NAME</u>
Alewife	ALW	<i>Alosa pseudoharengus</i>
American eel	EEL	<i>Anguilla rostrata</i>
American shad	ASD	<i>Alosa sapidissima</i>
American sand lance	ASL	<i>Ammodytes americanus</i>
Atlantic needlefish	ANF	<i>Strongylura marina</i>
Atlantic silversides	ASS	<i>Menidia menidia</i>
Atlantic tomcod	TOM	<i>Microgadus tomcod</i>
Banded butterflyfish	BBF	<i>Chaetodon striatus</i>
Banded gunnel	BGN	<i>Pholis fasciata</i>
Banded rudderfish	RUD	<i>Seriola zonata</i>
Bay anchovy	ACH	<i>Anchoa mitchilli</i>
Black-spot stickleback	BSS	<i>Gasterosteus wheatlandi</i>
Black sea bass	BSB	<i>Centropristes striata</i>
Blueback herring	BBH	<i>Alosa aestivalis</i>
Bluemass	BLF	<i>Pomatomus saltatrix</i>
Blue spotted coronetfish	BSC	<i>Fistularia tabacaria</i>
Crevalle jack	CRJ	<i>Caranx hippos</i>
Cunner	CUN	<i>Tautogolabrus adspersus</i>
Feather Blenny	FBL	<i>Hypsoblennius bentzi</i>
Flying Gurnard	FGD	<i>Dactylopterus volitans</i>
Four-spine stickleback	FSS	<i>Apeltes quadratus</i>
Gizzard Shad	GIZ	<i>Dorosoma cepedianum</i>
Gray snapper	GRA	<i>Lutjanus griseus</i>
Grubby	GRB	<i>Myoxocephalus aeneus</i>
Hogchoker	HOG	<i>Trinectes maculatus</i>
Inshore lizardfish	LIZ	<i>Synodens foetens</i>
Little skate	LSK	<i>Raja erinacea</i>
Menhaden	MEN	<i>Brevoortia tyrannus</i>
Moonfish	MOO	<i>Mene maculata</i>
Mummichog	MUM	<i>Fundulus heteroclitus</i>
Naked goby	NKG	<i>Gobiosoma boscii</i>
Nine-spine stickleback	NSS	<i>Pungitius pungitius</i>
Northern kingfish	NKF	<i>Menticirrhus saxatilis</i>
Northern pipefish	PIP	<i>Syngnathus fuscus</i>
Northern puffer	PUF	<i>Sphaerooides maculatus</i>
Northern searobin	NSR	<i>Prionotus carolinus</i>
Northern stargazer	STR	<i>Astroscopus guttatus</i>
Pumpkinseed	PUM	<i>Lepomis gibbosus</i>
Rainbow smelt	RSM	<i>Osmerus mordax</i>
Rainwater killifish	RWK	<i>Lucania parva</i>
Rock gunnel	RGN	<i>Pholis gunnellus</i>
Northern seahorse	SEH	<i>Hippocampus erectus</i>
Northern sennet	NOS	<i>Sphyraena borealis</i>
Scup	PGY	<i>Stenotomus chrysops</i>
Sheepshead minnow	SHM	<i>Cyprinodon variegatus</i>
Shorthorn Sculpin	SHS	<i>Myoxocephalus scorpius</i>
Skilletfish	SKL	<i>Gobiesox strumosus</i>
Smallmouth flounder	SMF	<i>Etropus microstomus</i>
Smooth dogfish	SMD	<i>Mustelus canis</i>
Spotted hake	SPH	<i>Urophycis regius</i>
Striped anchovy	STA	<i>Anchoa hepsetus</i>
Striped bass	STB	<i>Morone saxatilis</i>
Striped burrfish	SBF	<i>Chilomycterus schoepfi</i>
Striped killifish	SKF	<i>Fundulus majalis</i>
Striped searobin	SSR	<i>Prionotus evolans</i>
Summer flounder	SFL	<i>Paralichthys dentatus</i>
Tautog	BKF	<i>Tautoga onitis</i>
Three-spine stickleback	TSS	<i>Gasterosteus aculeatus</i>

Appendix 8.1, continued:

Toadfish	TDF	<i>Opsanus tau</i>
Weakfish	WKF	<i>Cynoscion regalis</i>
Web Burrfish	WBF	<i>Chilomycterus antillarum</i>
White mullet	WML	<i>Mugil curema</i>
Windowpane flounder	WPF	<i>Scophthalmus aquosus</i>
Winter flounder (YOY)	WFO	<i>Pseudopleuronectes americanus</i>
Winter flounder (AGE 1+)	WFL	<i>Pseudopleuronectes americanus</i>
Yellow jack	YJK	<i>Caranx bartholomaei</i>

Appendix 8.2: Invertebrate species captured in the Estuarine Seine Survey, 1988-2019.

<u>COMMON NAME</u>	<u>SPECIES CODE</u>	<u>SCIENTIFIC NAME</u>
Bay Scallop	SCA	<i>Argopecten irradians</i>
Blue crab	BCR	<i>Callinectes sapidus</i>
Brown Shrimp	BNS	<i>Panaeus aztecus</i>
Channeled Whelk	CHW	<i>Busycon canaliculatus</i>
Common Slipper Shell	CSL	<i>Crepidula fornicata</i>
Northern Comb Jelly	COM	<i>Bolinopsis infundibulum</i>
Green crab	GCR	<i>Carcinus maenas</i>
Flat-claw hermit crab	HER	<i>Pagurus spp.</i>
Horseshoe crab	HSC	<i>Limulus polyphemus</i>
Japanese crab	JSC	<i>Hemigrapsus sanguineus</i>
Jonah crab	JCR	<i>Cancer borealis</i>
Knobbed whelk	KNW	<i>Busycon carica</i>
Lady crab	LCR	<i>Ovalipes ocellatus</i>
Long-finned squid	SQI	<i>Loligo pealeii</i>
Mantis shrimp	MAN	<i>Squilla empusa</i>
Moon Jelly	MOJ	<i>Aurelia aurita</i>
Mud crab	BMC	<i>Panopeus spp.</i>
Mole crab	MLR	<i>Emerita talpoida</i>
Mud snail	MSN	<i>Nassarius obsoletus</i>
Northern moon snail	NMS	<i>Lunatia heros</i>
Oyster drill	OYD	<i>Urosalpinx cinerea</i>
Rock crab	RCR	<i>Cancer irroratus</i>
Sand shrimp	CRG	<i>Crangon septemspinosa</i>
Sea Star	STF	<i>Asterias forbesi</i>
Shore shrimp	PAL	<i>Palaemonetes spp.</i>
Spider crab	SPI	<i>Libinia emarginata</i>